

ROAD TRAFFIC NOISE IN URBAN ENVIRONMENTS IN AHVAZ CITY, IRAN

Mohammad Javad Mohammadi^{1,2}, Esmail Charkhloo³, Sahar Geravandi¹, Afshin Takdastan⁴, Somayeh Rahimi¹, Ahmad Reza Yari⁵, Rashin Alizadeh⁶, Mehdi Vosoughi², Sina Dobaradaran⁷, Parviz Javanmardi^{8,*}

¹Abadan school of Medical Sciences, Abadan, Iran

²Student Research Committee, Department of Environmental Health Engineering, School of Public Health AND Environmental Technologies Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

³Department of Environmental Health Engineering, Faculty of Health, Jiroft University of Medical Sciences, Jiroft, Iran

⁴Environmental Technologies Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁵Research Center for Environmental Pollutants, Qom University of Medical Sciences, Qom, Iran

⁶Department of Health Education and Promotion, School of Health, Tehran University of Medical Sciences, Tehran, Iran

⁷Department of Environmental Health Engineering, Faculty of Health & the Persian Gulf Marine Biotechnology Research Center, the Persian Gulf Research Center, Bushehr University of Medical Sciences, Bushehr, Iran.

⁸Health Care System of Hendejan, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

ABSTRACT

Noise pollution because of physiological and psychological effects on human is one of the importance environmental pollutants. The most important sources of this pollution are transportation and car traffic. The present study was implemented with an overall goal of assessment of noise pollution from traffic in Ahvaz, Iran in 2012. In this descriptive study, 7 roads of the main routes of Ahvaz which have high density of traffic are selected and 134 measuring stations were chosen. At measuring stations, 4 days of week, three time period, and 1523 measurements were recorded totally. Each measurement was recorded for 30 minutes. After transferring data to computer, SPSS and Excel software were applied for statistical analysis; also T-test and ANOVA were used for statistical tests. According to result this study the average of equivalent sound pressure level in all stations is equal to 76.28 ± 3.12 dB. There was a statistically significant difference between the mean values of equivalent sound pressure level on different roads, different weekdays and various daytimes ($P < 0.05$). According to the results, it was found that Ahvaz has a higher noise pollution in comparison with outdoor permissible limit standard in Iran. So, it should seriously be taken into consideration in order to control noise and prevent the effects.

KEYWORDS:

Noise pollution, Noise Index, Traffic, Ahvaz, Iran.

INTRODUCTION

Developing industrial cities and tendency to live in large cities have caused people in metropolises face a number of problems [1-4]. Among the problems that can endanger human serenity and health is noise pollution [5-7]. According result different studies number of person's exposure to railway, aircraft and industry noise affects are 16, 4 and 1 million persons [5-7]. Exposure to road traffic noise can be causes numerous adverse health endpoints [8-13]. The main sources of noise pollution are traffic and transportation [14-16]. The capital of Khuzestan province with a population of 1.2 million persons and area of 140 square kilometers is Ahvaz city [17-20]. Significant urban and industrial developments and the growth of universities have increased transportation on roads and routes so that the total number of vehicles (including bicycles, motorcycles, talkies, cars, trucks, heavy vehicles, etc.) is over 336710, which caused different kinds of pollution especially noise pollution in different parts of the city [21]. In megacities at developed country one of most important major source of community noise, especially near a highways is the road traffic. Result of several studies demonstrated the impact of this highway and road nuisance [22,23]. In the Gomez article, noise pollutions from traffic were discussed as a factor of society health level reduction [24]. Also, in another study in Zanjan, Iran L_{10} , L_{50} , L_{90} , SPL and NPL indexes are evaluated and showed a noise level above the standard in different parts of the city [25]. Another study by Rahmani et al. in 2011 evaluated noise level and developed two noise determinations, which showed noise level above




FIGURE 1

Mapping of Ahvaz with location of measuring stations

standard in Iran [26]. In another study, Lee and et al, evaluated the model road traffic noise levels and estimate the human exposure at 25 districts in the metropolitan Seoul, Republic of Korea [27]. Pichai analyzed and assessed the effect of traffic noise at motorway in newly expanded Bangkok-Chonburi [28]. Another study to evaluate noise levels in Jordan showed that influential factors on noise levels are traffic density, speed, honking, and number of lines, line widths, road slope and pavement [29]. Gundoghu et al showed that the average and maximum day noise level in Erzurum, Turkey was above standard [30]. The results of a study modeling of Road traffic noise in Brazil in 2003 indicated that people who live or work nearby these roads, experience above standard noise levels [31]. Also the result of a study conducted by Ali and Tamura in Cairo, Egypt, showed above the standard noise level [32]. One cross-sectional study has found a higher but non-significant prevalence of hypertension among male workers exposed to noise levels at frequencies of 2000, 4000 and 8000 Hz [33].

The main purpose of this study was the evaluation of noise pollution from the traffic of Ahvaz, Iran during the year 2012 and comparing the results with standards.

MATERIALS AND METHODS

Study population and exposure assessment.

The descriptive study assessed noise index on 7 city roads (Pasdaran and Ayatollah – Behbahani, Enghelab, shahid–Chamran and Golestan, Shariati and Azadegan) in 134 given stations in summer 2012. Squares, intersections and spots between them on these roads were under study [34-36]. Measurements were done in 4 days (Sat, Tues, Thurs, Fri) representing workdays and weekends [34]. According Environmental Protection Agency (EPA), samples were mostly gathered at rush hours based on investigations, 7-8 AM, 12-13 PM and 8-9

PM. [37,38]. Noise levels were determined by CEL/450 scales which are accurate and made based on international standards. This device has various models and many applications, so it can measure several variables [25]. The scale was calibrated before use and when it was stopped, protective foam was used to control air effect on the microphone [39,40]. Measuring was done without catching people, pedestramts, drivers, shop keepers or landowner's attention. The scale was set at the height of 1.6m, in the position dB fast and sampling was done in 30 minutes [41-43]. All measurements were done on both sides of the roads [42,43]. The studied indexes are as follows; L_{eq} (Sound pressure Level), L_{max} (Maximum sound Pressure Level), L_{min} (Minimum sound Pressure Level) [39].

Statistical Analysis. In the present study, we used descriptive statistics (frequency, mean, and standard deviation for each variable). The data were analyzed by Excel, SPSS, ANOVA and T- test.

RESULTS AND DISCUSSION

The results showed that the average of equivalent sound level (L_{eq}) in all stations was 76.28 ± 3.08 dB, which was higher than the environmental standard in EPA and Iran (65dB) [44]. Also in all studied roads, days and time periods, noise level was above the standard. The minimum and maximum sound pressure Levels were 76.28 ± 3.18 and 68.93 ± 3.36 respectively. Pasdaran and Behbahani Highways with 73.65 ± 6.20 and 73.09 ± 4.20 , and Shahid–chamran and Azadegan highways with 71.83 ± 2.05 and 71.21 ± 6.43 showed the highest and lowest average equivalent sound pressure levels, respectively. Tables 1-2 and Figure 2 show noise pollution parameters of the present study.

Also by ANOVA and T-test, there was a considerable difference between the average of

equivalent sound level at different roads, days, and time periods, on Friday and other days of the week ($P < 0.05$). But at different stations (on squares, intersections and interval spots) no significant difference was observed between L_{eq} .

TABLE 1
The results of measured sound parameters in Ahvaz, Iran. (2012)

Parameter Value	L_{max}	L_{min}	L_{eq}
Min	69.12	57.25	61.21
Max	95.46	80.62	91.35
Average	84.32	68.93	76.28

TABLE 2
The average of noise pollution parameters for the studied times in Ahvaz, Iran (2012)

Parameter Hours	L_{max}	L_{min}	L_{eq}
6-8 A.M	83.19	83.19	72.09
12-13 P.M	83.11	83.11	72.44
8-9 P.M	84.02	84.02	73.34




FIGURE 2
The average of Equivalent Sound Pressure Level in the studied roads

The results showed that the average of equivalent sound level at all stations was above standard (65dB). Some studies reported similar equivalent levels for traffic noise range. According to report of Babisch in 2014 there was a significant relationship between coronary heart diseases and road traffic noise. They demonstrated that per increase of the weighted day-night noise level L_{DN} of 10 dB (A) can be increase of the relative risk of 1.08 (95% confidence interval: 1.04, 1.13) [44]. Golmohammadi (2005-2006) reported that the value for Hamadan, Iran, in all stations to be 69.04 ± 4.25 (dB) [45]. Motallebi Kashani et al in Kashan, Iran, (2000- 2001) reported that the crowded spots hospital rooms, and residential – industrial areas were 76.7, 65.6, 56.7, 58.7 dB respectively [46]. Belojevic et al. (1977) reported that the noise level in crowded areas of downtown in Belgrad was above 65 dB (daytime equivalent level 76.5 and the night were 64.5 dB) [47]. A study conducted in Messina, Italy showed L_{eq} above 75 dB [48]. Another study reported the traffic noise level in more than 60

crowded areas in 8 cities of Nigeria by momentary sampling 24 hrs. Equivalent sound level and Maximum environmental sound levels were 84.6 and 1.5 dB in Net A respectively. High maximum environmental sound level was considered, pertain to cultural factors, including big African horns by motorcycles [49]. Alizadeh et al reported the average sound pressure indexes (L_{10} – L_{50} , L_{90} , LD) in measuring stations in Sarri, Iran, above standard (65dB) [50]. The considerable difference between the average equivalent sound level at different roads, days, time periods, on Friday and other days of the week ($P < 0.05$), owing to the passing vehicles load and their speed mainly. Naddafi and Golmohammadi reported the same results [39,45]. Noise pollution in big cities is still a matter of serious social concern; some of whose causes could be construction and traffic engineering ones. According to the results, Noise pollution in Ahvaz, Iran is due to the old urban residential area traffic issues significantly.

CONCLUSION

According to the results, it was found that Ahvaz has a higher noise pollution in comparison with outdoor permissible limit standard in Iran. So noise pollution is a matter of serious concern and causes need to be studied. Traffic management reforms, using noise controllers, making one way streets, planting on sides of the roads to absorb the noise, locating commercial centers out of residential areas, discarding old vehicles, removing bumps, using high speed reducers such as traffic signs, speed limits and speed bumps on lowly traveled streets in order to reduce noise pollution in different areas of the city seems to be necessary.

ACKNOWLEDGEMENT

The authors would take to thank the Azma jonob company for their technical support and providing the facilities.

Conflict of Interests. Authors have no conflict of interests.

Authors Contributions. Study concept, design and critical revision of the manuscript for important intellectual content: Mohammad Javad Mohammadi, Sahar Geravandi, Esmail Charkhloo, Afshin Takdastan, Ahmad Reza Yari, Mohammad Mahboubi, Ghollam Abas Shirali, Sina Dobaradaran, Somayeh Rahimi, Mehdi Vosoughi, Rashin Alizadeh, drafting of the manuscript and advisor. Drafting of the manuscript and advisor; doing experiments Parviz Javanmardi.

Financial Disclosure. We have no financial

interests related to the material in the manuscript.

Funding/Support. This work was financially supported by Ahvaz Jundishapur University of Medical Sciences.

REFERENCES

- [1] Davar H, Taghavirad SS, Mohammadi MJ (2014) The investigation of effects of silica on the environment and prevention of release of the silica particles with simulation of gas-solid flow in a gas cyclone *Research Journal of Chemistry and Environment* 18 (11):28-30
- [2] Dobaradaran S, Geravandi S, Goudarzi G, Idani E, Salmanzadeh S, Soltani F, Yari AR, Mohammadi MJ (2016) Determination of Cardiovascular and Respiratory Diseases Caused by PM10 Exposure in Bushehr, 2013. *Journal of Mazandaran University of Medical Sciences* 26 (139):42-52
- [3] Goudarzi G, Geravandi S, Idani E, Hosseini SA, Baneshi MM, Yari AR, Vosoughi M, Dobaradaran S, Shiralil S, Marzooni MB (2016) An evaluation of hospital admission respiratory disease attributed to sulfur dioxide ambient concentration in Ahvaz from 2011 through 2013. *Environmental Science and Pollution Research* 23 (21):22001-22007
- [4] Khaefi M, Goudarzi G, Yari AR, Geravandi S, Dobaradaran S, Idani E, Javanmardi P, Yousefi F, Hashemzadeh B, Shahriari A, Javad MM (2016) An association between ambient pollutants and hospital admitted respiratory cases in Ahvaz, Iran. *Fresen. Environ. Bull.* 25 (10):3955-3961
- [5] Rösli M (2013) [Health effects of environmental noise exposure]. *Therapeutische Umschau Revue thérapeutique* 70 (12):720-724
- [6] Sørensen M, Lühndorf P, Ketzel M, Andersen ZJ, Tjønnelund A, Overvad K, Raaschou-Nielsen O (2014) Combined effects of road traffic noise and ambient air pollution in relation to risk for stroke? *Environmental research* 133:49-55
- [7] Sarnat JA, Golan R, Greenwald R, Raysoni AU, Kewada P, Winkquist A, Sarnat SE, Flanders WD, Mirabelli MC, Zora JE (2014) Exposure to traffic pollution, acute inflammation and autonomic response in a panel of car commuters. *Environmental research* 133:66-76
- [8] Chang T-Y, Beelen R, Li S-F, Chen T-I, Lin Y-J, Bao B-Y, Liu C-S (2014) Road traffic noise frequency and prevalent hypertension in Taichung, Taiwan: a cross-sectional study. *Environmental Health* 13 (1):1
- [9] Sørensen M, Andersen ZJ, Nordsborg RB, Jensen SS, Lillilund KG, Beelen R, Schmidt EB, Tjønnelund A, Overvad K, Raaschou-Nielsen O (2012) Road traffic noise and incident myocardial infarction: a prospective cohort study. *PLoS One* 7 (6):e39283
- [10] Selander J, Bluhm G, Nilsson M, Hallqvist J, Theorell T, Willix P, Pershagen G (2013) Joint effects of job strain and road-traffic and occupational noise on myocardial infarction. *Scandinavian journal of work, environment & health*:195-203
- [11] Bluhm GL, Berglund N, Nordling E, Rosenlund M (2007) Road traffic noise and hypertension. *Occupational and environmental medicine* 64 (2):122-126
- [12] Van Kempen E, Babisch W (2012) The quantitative relationship between road traffic noise and hypertension: a meta-analysis. *Journal of hypertension* 30 (6):1075-1086
- [13] Barregard L, Bonde E, Öhrström E (2009) Risk of hypertension from exposure to road traffic noise in a population-based sample. *Occupational and environmental medicine* 66 (6):410-415
- [14] Bodin T, Albin M, Ardö J, Stroh E, Östergren P-O, Björk J (2009) Road traffic noise and hypertension: results from a cross-sectional public health survey in southern Sweden. *Environmental Health* 8 (1):1
- [15] Chang T-Y, Liu C-S, Bao B-Y, Li S-F, Chen T-I, Lin Y-J (2011) Characterization of road traffic noise exposure and prevalence of hypertension in central Taiwan. *Science of the total environment* 409 (6):1053-1057
- [16] Istaito T, Houthuijs D, Lebret E (2014) Multi-country willingness to pay study on road-traffic environmental health effects: are people willing and able to provide a number? *Environmental Health* 13 (1):1
- [17] Geravandi S, Goudarzi GR, Vosoughi Niri M, Mohammadi Mj, Saeidimehr S, Geravandi S (2015) Estimation of the cardiovascular and respiratory mortality rate resulted from exposure to sulfur dioxide pollutant in ahvaz. *Journal of Environmental Studies* 41 (2):341-350
- [18] Goudarzi G, Geravandi S, Forouzandeh H, Babaei AA, Alavi N, Niri MV, Khodayar MJ, Salmanzadeh S, Mohammadi MJ (2015) Cardiovascular and respiratory mortality attributed to ground-level ozone in Ahvaz, Iran. *Environmental monitoring and assessment* 187 (8):1-9
- [19] Neisi A, Goudarzi G, Akbar Babaei A, Vosoughi M, Hashemzadeh H, Naimabadi A, Mohammadi MJ, Hashemzadeh B (2016) Study of heavy metal levels in indoor dust and their health risk assessment in children of Ahvaz city, Iran. *Toxin Reviews* 35 (1-2):16-23
- [20] Yari AR, Goudarzi G, Geravandi S, Dobaradaran S, Yousefi F, Idani E, Jamshidi F, Shiralil S, Khishdost M, Mohammadi MJ (2016) Study of ground-level ozone and its health risk

- assessment in residents in Ahvaz City, Iran during 2013. *Toxin Reviews* 35 ((3–4)):201–206
- [21] Nadim F, Bagtzoglou AC, Afshar A, Iranmahboob J, Monazzam MR, Yunesian M (2009) Transport Management in Iran: Traffic Safety and Cleaner Cars. *Journal of Environmental Engineering and Management* 19 (6):317–325
- [22] Ismail A, Mansor MRA, Nor MJM, Nuawi MZ, Zulkifli R, Rahman M, Sopian K (2008) Comparative studies and impact analysis of environmental noise modeling from Malaysian industrial projects. *WSEAS Trans Environ Dev* 4:666–675
- [23] Parbat D, Nagarnaik P (2006) Assessment of Traffic Noise Levels and its Annoyance on Arterial Roads of Intermediate City. *WSEAS Transactions on Environment and Development* 12 (2):1443–1448
- [24] Gómez-Jacinto L, Moral-Toranzo F (1999) Urban traffic noise and self-reported health. *Psychological reports* 84 (3 suppl):1105–1108
- [25] Naddafi K, Yunesian M, Mesdaghinia A, Mahvi A, Asgari A (2008) Noise pollution in Zanjan city in 2007. *ZUMS Journal* 16 (62):85–96
- [26] Rahmani S, Mousavi SM, Kamali M (2011) Modeling of road-traffic noise with the use of genetic algorithm. *Applied Soft Computing* 11 (1):1008–1013
- [27] Lee J, Gu J, Park H, Yun H, Kim S, Lee W, Han J, Cha J-S (2014) Estimation of populations exposed to road traffic noise in districts of Seoul metropolitan area of Korea. *International journal of environmental research and public health* 11 (3):2729–2740
- [28] Pamanikabud P (2009) Impact Assessment of Motorway Traffic Noise Using Visualized Noise Mapping Technique. *WSEAS Trans Environ Dev* 5:24–33
- [29] Abo-Qudais S, Alhiary A (2007) Statistical models for traffic noise at signalized intersections. *Building and Environment* 42 (8):2939–2948
- [30] Gündoğdu Ö, Gökdağ M, Yüksel F (2005) A traffic noise prediction method based on vehicle composition using genetic algorithms. *Applied Acoustics* 66 (7):799–809
- [31] Calixto A, Diniz FB, Zannin PH (2003) The statistical modeling of road traffic noise in an urban setting. *Cities* 20 (1):23–29
- [32] Ali SA, Tamura A (2003) Road traffic noise levels, restrictions and annoyance in Greater Cairo, Egypt. *Applied Acoustics* 64 (8):815–823
- [33] Chang T-Y, Liu C-S, Young L-H, Wang V-S, Jian S-E, Bao B-Y (2012) Noise frequency components and the prevalence of hypertension in workers. *Science of the total environment* 416:89–96
- [34] Wetzel E, Nicolas J, Andre P, Boreux J-J (1999) Modelling the propagation pathway of street-traffic noise: practical comparison of German guidelines and real-world measurements. *Applied Acoustics* 57 (2):97–107
- [35] Nejadkoorki F, Yousefi E, Naseri F (2010) Analysing street traffic noise pollution in the city of Yazd. *Journal of Environmental Health Science & Engineering* 7 (1):53–62
- [36] Geravandi S, Takdastan A, Zallaghi E, Niri MV, Mohammadi MJ, Saki H, Naiemabadi A (2015) Noise pollution and health effects. *Jundishapur Journal of Health Sciences* 7 (1)
- [37] Hogan CM (1973) Analysis of highway noise. *Water, Air, and Soil Pollution* 2 (3):387–392
- [38] Bluhm G, Nordling E, Berglind N (2004) Road traffic noise and annoyance-An increasing environmental health problem. *Noise and Health* 6 (24):43
- [39] Golmohammadi R, Aliabadi M (2012) Noise Pollution and its Irritating Effects in Hospitals of Hamadan, Iran. *Health System Research* 7 (6):958–964
- [40] Fidell S (1978) Nationwide urban noise survey. *The Journal of the Acoustical Society of America* 64 (1):198–206
- [41] Ranjbar HR, Gharagozlou AR, Nejad ARV (2012) 3D analysis and investigation of traffic noise impact from Hemmat Highway located in Tehran on buildings and surrounding areas. *Journal of Geographic Information System* 4 (4):322
- [42] Tansatcha M, Pamanikabud P, Brown A, Affum JK (2005) Motorway noise modelling based on perpendicular propagation analysis of traffic noise. *Applied acoustics* 66 (10):1135–1150
- [43] Malakootian M, Ahmadian M, Yaghmaeian K, Dowlatshahi S, Ravandi MG (2012) Level Changes of Traffic Noise in Kerman City, Southeast Iran. *Iranian journal of public health* 41 (1):107
- [44] Babisch W (2014) Updated exposure-response relationship between road traffic noise and coronary heart diseases: A meta-analysis. *Noise and Health* 16 (68):1–9.
- [45] R G (2007) Develop and design a traffic noise model for cities Main roads (case study in Hamedan) Islamic Azad university, science and Research Branch of Tehran PHD thesis:87–98
- [46] Mottalebi kashani M HM, Akbari H, Almasi H (2002) Assessment of noise pollution value in kashan city in 2000 – 2001. *feyz scientific* 21:30–32
- [47] Belojević G, Jakovljević B, Aleksić O (1997) Subjective reactions to traffic noise with regard to some personality traits. *Environment International* 23 (2):221–226
- [48] Piccolo A, Plutino D, Cannistraro G (2005) Evaluation and analysis of the environmental noise of Messina, Italy. *Applied Acoustics* 66 (4):447–465

- [49] Onuu M (2000) Road traffic noise in Nigeria: measurements, analysis and evaluation of nuisance. *Journal of Sound and Vibration* 233 (3):391-405
- [50] Ali zadeh A MM, Eatemadi Nejad S, Yazdani cherati J (2009) Assessment of noise pollution in Sarri during one year (2007-2008). *Jurnal of mazandaran medical science university* 19 (62):45-52

Received: 03.12.2016

Accepted: 10.03.2017

CORRESPONDING AUTHOR

Parviz Javanmardi

Health Care System of Hendejan, Ahvaz
Jundishapur University of Medical Sciences, Ahvaz,
Iran

e-mail: parvizjavanmardi1394@gmail.com